



ELSEVIER

Available online at www.sciencedirect.com

ScienceDirect

Linear Algebra and its Applications 428 (2008) 1791–1802

LINEAR ALGEBRA
AND ITS
APPLICATIONSwww.elsevier.com/locate/laa

Property (w) for perturbations of polaroid operators[☆]

Pietro Aiena^{a,*}, Jesús R. Guillen^b, Pedro Peña^c^a *Dipartimento di Metodi e Modelli Matematici, Facoltà di Ingegneria, Università di Palermo, Viale delle Scienze, I-90128 Palermo, Italy*^b *Departamento de Matemáticas, Facultad de Ciencias, Universidad ULA, Merida, Venezuela*^c *Departamento de Física y Matemática, NURR, ULA-Trujillo, Venezuela*

Received 13 December 2006; accepted 16 October 2007

Submitted by V. Mehrmann

Abstract

A bounded linear operator $T \in L(X)$ acting on a Banach space satisfies property (w) , a variant of Weyl's theorem, if the complement in the approximate-point spectrum $\sigma_a(T)$ of the Weyl essential approximate-point spectrum $\sigma_{wa}(T)$ is the set of all isolated points of the spectrum which are eigenvalues of finite multiplicity. In this note, we study the stability of property (w) for a polaroid operator T acting on a Banach space, under perturbations by finite rank operators, by nilpotent operators and, more generally, by algebraic operators commuting with T .

© 2007 Elsevier Inc. All rights reserved.

AMS classification: Primary 47A10, 47A11; Secondary 47A53, 47A55*Keywords:* Property (w) ; Weyl's theorems; Polaroid operators

1. Definitions and basic results

Let X denote an infinite-dimensional complex Banach space, and denote by $L(X)$ the algebra of all bounded linear operators on X . If $T \in L(X)$ by $\alpha(T)$ we shall denote the dimension of the kernel $\ker T$ and by $\beta(T)$ the codimension of the range $T(X)$. Recall that $T \in L(X)$ is said to be *upper semi-Fredholm*, $T \in \Phi_+(X)$, if $T(X)$ is closed and $\alpha(T) < \infty$, while $T \in L(X)$ is called

[☆] This research was supported by Fondi ex-60, 2006, Università di Palermo.

* Corresponding author.

E-mail addresses: paiana@unipa.it (P. Aiena), rguillen@ula.ve (J.R. Guillen), pedrop@ula.ve (P. Peña).

lower semi-Fredholm, $T \in \Phi_-(X)$, if $\beta(T) < \infty$. The class of all semi-Fredholm operators is defined by $\Phi_{\pm}(X) := \Phi_+(X) \cup \Phi_-(X)$, while the class of all Fredholm operators is defined by $\Phi(X) := \Phi_+(X) \cap \Phi_-(X)$. If $T \in \Phi_{\pm}(X)$, the *index* of T is defined by $\text{ind}(T) := \alpha(T) - \beta(T)$. Define

$$W_+(X) := \{T \in \Phi_+(X) : \text{ind } T \leq 0\},$$

and

$$W_-(X) := \{T \in \Phi_-(X) : \text{ind } T \geq 0\}.$$

The set of *Weyl operators* is defined by

$$W(X) := W_+(X) \cap W_-(X) = \{T \in \Phi(X) : \text{ind } T = 0\}.$$

The classes of operators defined above generate the following spectra. The *Weyl spectrum* is defined by

$$\sigma_w(T) := \{\lambda \in \mathbb{C} : \lambda I - T \notin W(X)\},$$

while the *Weyl essential approximate-point spectrum* is defined by

$$\sigma_{wa}(T) := \{\lambda \in \mathbb{C} : \lambda I - T \notin W_+(X)\},$$

and analogously, the *Weyl essential surjectivity spectrum* is defined by

$$\sigma_{ws}(T) := \{\lambda \in \mathbb{C} : \lambda I - T \notin W_-(X)\},$$

Note that $\sigma_w(T) = \sigma_{wa}(T) \cup \sigma_{ws}(T)$. By duality we also have $\sigma_{wa}(T) = \sigma_{ws}(T^*)$ and $\sigma_{wa}(T^*) = \sigma_{ws}(T)$. The classical *approximate-point spectrum* is defined by

$$\sigma_a(T) := \{\lambda \in \mathbb{C} : \lambda I - T \text{ is not injective or has not closed range}\},$$

while the *surjective spectrum* is defined by

$$\sigma_s(T) := \{\lambda \in \mathbb{C} : \lambda I - T \text{ is not onto}\}.$$

It is well-known that $\sigma_a(T^*) = \sigma_s(T)$ and $\sigma_s(T^*) = \sigma_a(T)$. The *ascent* of $T \in L(X)$ is defined as $p := p(T) = \inf\{n \in \mathbb{N} : \ker T^n = \ker T^{n+1}\}$, while the *descent* is defined as $q := q(T) = \inf\{n \in \mathbb{N} : T^n(X) = T^{n+1}(X)\}$, where the infimum over the empty set is taken ∞ . It is well-known that if $p(T)$ and $q(T)$ are both finite then $p(T) = q(T)$ [14, Proposition 38.3]. Moreover, $0 < p(\lambda I - T) = q(\lambda I - T) < \infty$ precisely when λ is a pole of the resolvent of T , see [14, Proposition 50.2].

The class of all *Browder operators* is defined

$$B(X) := \{T \in \Phi(X) : p(T) = q(T) < \infty\}.$$

By Theorem 3.4 of [1] we have $B(X) \subseteq W(X)$. The *Browder spectrum* of $T \in L(X)$ is defined by

$$\sigma_b(T) := \{\lambda \in \mathbb{C} : \lambda I - T \notin B(X)\}.$$

Clearly, $\sigma_w(T) \subseteq \sigma_b(T)$ for all $T \in L(X)$, and if $\lambda \in \sigma(T) \setminus \sigma_b(T)$ then λ is isolated in $\sigma(T)$. For a bounded operator $T \in L(X)$, define

$$\pi_{00}(T) := \{\lambda \in \text{iso } \sigma(T) : 0 < \alpha(\lambda I - T) < \infty\},$$

and

$$\pi_{00}^a(T) := \{\lambda \in \text{iso } \sigma_a(T) : 0 < \alpha(\lambda I - T) < \infty\}.$$

Following Coburn [9], we shall say that $T \in L(X)$ satisfies *Weyl's theorem* if

$$\sigma(T) \setminus \sigma_w(T) = \pi_{00}(T). \quad (1)$$

The following two interesting variants of Weyl's theorem have been introduced by Rakočević [20,21] and studied in the recent paper [6].

Definition 1.1. A bounded operator $T \in L(X)$ is said to satisfy property (w) if

$$\sigma_a(T) \setminus \sigma_{wa}(T) = \pi_{00}(T),$$

while $T \in L(X)$ is said to satisfy a -Weyl's theorem if

$$\sigma_a(T) \setminus \sigma_{wa}(T) = \pi_{00}^a(T).$$

As observed in [6], we have either of a -Weyl's theorem or property (w) for $T \Rightarrow$ Weyl's theorem holds for T .

Examples of operators satisfying Weyl's theorem but not property (w) may be found in [6]. Property (w) is independent from a -Weyl's theorem: In [6] there are examples of operators $T \in L(X)$ satisfying property (w) but not a -Weyl's theorem and vice versa. Note that property (w) is fulfilled by a relevant number of Hilbert space operators, see [6]. For instance, property (w) is satisfied by generalized scalar operators, and whenever the Hilbert space adjoint T' has property $H(p)$ [6, Corollary 2.20] (the terminology is defined in the next section). Generally, property (w) , as well as Weyl's theorems, do not survive under perturbations. More can be said: Weyl's theorems and property (w) for a bounded operator T are liable to fail also under “small” perturbations K , if “small” is interpreted in the sense of compact or quasi-nilpotent operators. In [3] some sufficient conditions are given for which we have the stability of property (w) , under perturbations by finite rank operators, compact operators, or quasi-nilpotent operator commuting with T . In this paper we shall prove that the stability of property (w) holds whenever T belongs to some special classes of operators and K is a commuting algebraic operator that commutes with T .

2. Results

Recall that the operator $T \in L(X)$ is said to have *the single valued extension property* at $\lambda_0 \in \mathbb{C}$ (abbreviated SVEP at λ_0), if for every open disc \mathbb{D} centered at λ_0 , the only analytic function $f : \mathbb{D} \rightarrow X$ which satisfies the equation $(\lambda I - T)f(\lambda) = 0$ for all $\lambda \in \mathbb{D}$ is the function $f \equiv 0$. An operator $T \in L(X)$ is said to have SVEP if T has SVEP at every point $\lambda \in \mathbb{C}$.

Clearly, an operator $T \in L(X)$ has SVEP at every point of the resolvent $\rho(T) := \mathbb{C} \setminus \sigma(T)$. Moreover, the identity theorem for analytic functions implies that $T \in L(X)$ has SVEP at every point of the boundary $\partial\sigma(T)$ of the spectrum $\sigma(T)$. In particular, T has SVEP at every isolated point of the spectrum. Note that

$$p(\lambda I - T) < \infty \Rightarrow T \text{ has SVEP at } \lambda, \quad (2)$$

dually

$$q(\lambda I - T) < \infty \Rightarrow T^* \text{ has SVEP at } \lambda. \quad (3)$$

Also

$$\sigma_a(T) \text{ does not cluster at } \lambda \Rightarrow T \text{ has SVEP at } \lambda. \quad (4)$$

Remark 2.1. The implications (2), (3) and (4) are equivalences whenever $\lambda I - T \in \Phi_{\pm}(X)$, see [1, Theorem 3.16].

Theorem 2.2. [6] *Let $T \in L(X)$. Then the following equivalences hold:*

- (i) *If T^* has SVEP, then property (w), Weyl's theorem, and a -Weyl's theorem for T are equivalent.*
- (ii) *If T has SVEP, then property (w), Weyl's theorem and a -Weyl's theorem for T^* are equivalent.*

Remark 2.3. In the case of Hilbert space operators $T \in L(H)$ instead of the dual T^* it is more appropriate to consider the Hilbert space adjoint T' . It should be noted that T^* has SVEP is equivalent to saying that T' has SVEP, see [2, Section 4].

The following result will be useful in the sequel.

Theorem 2.4. *Suppose that $T \in L(H)$, H a Hilbert space. Then property (w) holds for T^* if and only if property (w) holds for T' .*

Proof. (i) By means of the classical Fréchet–Riesz representation theorem we know that if U is the conjugate-linear isometry that associates to each $y \in H$ the linear form $x \rightarrow \langle x, y \rangle$ then

$$\bar{\lambda}I - T' = (\lambda I - T)' = U^{-1}(\lambda I - T)^*U. \quad (5)$$

This obviously implies that $\alpha(\bar{\lambda}I - T') = \alpha(\lambda I^* - T^*)$ for all $\lambda \in \mathbb{C}$. Since $\sigma(T') = \overline{\sigma(T)} = \overline{\sigma(T^*)}$ and analogously $\sigma_a(T') = \overline{\sigma_a(T^*)}$, it then easily follows that the equality $\pi_{00}(T') = \overline{\pi_{00}(T^*)}$ holds.

Suppose now that T^* satisfies property (w). From (5) we also have $\sigma_{aw}(T') = \overline{\sigma_{aw}(T^*)}$, so

$$\begin{aligned} \sigma_a(T') \setminus \sigma_{aw}(T') &= \overline{\sigma_a(T^*)} \setminus \overline{\sigma_{aw}(T^*)} = \overline{\sigma_a(T^*) \setminus \sigma_{aw}(T^*)} \\ &= \overline{\pi_{00}(T^*)} = \pi_{00}(T'), \end{aligned}$$

so T' satisfies property (w). The opposite implication follows in a similar way. \square

A bounded operator $T \in L(X)$ is said to be *polaroid* if $\text{iso } \sigma(T) = \emptyset$ or every isolated point of $\sigma(T)$ is a pole of the resolvent of T . Weyl's theorem for polaroid operators has been studied in a very recent paper by Duggal [11].

Theorem 2.5. *If $T \in L(X)$, X a Banach space, is polaroid then its dual T^* is polaroid. If $T \in L(H)$, H a Hilbert space, then T is polaroid if and only if its Hilbert space adjoint T' is polaroid.*

Proof. Suppose that $\lambda \in \text{iso } \sigma(T^*)$. Since $\sigma(T) = \sigma(T^*)$ then λ is an isolated point of $\sigma(T)$ so, by assumption, a pole of the resolvent of T . Let $p := p(\lambda I - T) = q(\lambda I - T)$. By Theorem 3.6 of [1] then

$$X = \ker(\lambda I - T)^p \oplus (\lambda I - T)^p(X). \quad (6)$$

Denote by M^{\perp} the annihilator of a subset M of X . Note that $(\lambda I - T)^p(X)$ is closed, since it is the kernel of the spectral projection associated with the spectral set $\{\lambda\}$, see Theorem 3.74 of

[1], hence by the classical closed range theorem $(\lambda I - T)^p(X)^\perp = \ker(\lambda I^* - T^*)^p$. From the decomposition (6) we then obtain

$$X^* = \ker(\lambda I - T)^{p\perp} \oplus (\lambda I - T)^p(X)^\perp = (\lambda I^* - T^*)^p(X^*) \oplus \ker(\lambda I^* - T^*)^p,$$

and hence, again by [1, Theorem 3.6], $p(\lambda I^* - T^*) = q(\lambda I^* - T^*) < \infty$, thus λ is a pole of the resolvent of T^* .

Suppose that $T \in L(H)$ is polaroid. To prove that T' is polaroid let λ be an isolated point in $\sigma(T') = \overline{\sigma(T)}$. Then $\bar{\lambda}$ is isolated in $\sigma(T)$, and hence $\bar{\lambda}$ is a pole of the resolvent of T , thus $p := p(\bar{\lambda}I - T) = q(\bar{\lambda}I - T) < \infty$. Consequently, $H = \ker(\bar{\lambda}I - T)^p \oplus (\bar{\lambda}I - T)^p(X)$ and the range $(\bar{\lambda}I - T)^p(X)$ is closed. From this it then follows that

$$H = (\ker \bar{\lambda}I - T)^{p\perp} \oplus (\bar{\lambda}I - T)^p(H)^\perp = (\lambda I - T')^p(H) \oplus \ker(\lambda I - T')^p,$$

where here N^\perp denotes the orthogonal complement of $N \subseteq H$. Therefore $p(\lambda I - T') = q(\lambda I - T') < \infty$, or equivalently λ is a pole of the resolvent of T' , thus T' is polaroid. Conversely, if T' is polaroid by the first part of the proof then $T'' = T$ is polaroid. \square

Theorem 2.6. Suppose that $T \in L(X)$.

- (i) If T^* has SVEP then $\sigma_{\text{wa}}(T) = \sigma_b(T)$.
- (ii) If T has SVEP then $\sigma_{\text{wa}}(T^*) = \sigma_b(T)$.

Proof. (i) The inclusions $\sigma_{\text{wa}}(T) \subseteq \sigma_w(T) \subseteq \sigma_b(T)$ hold for every $T \in L(X)$. Suppose that $\lambda \notin \sigma_{\text{wa}}(T)$. Then $\lambda I - T \in \Phi_+(X)$ with $\text{ind}(\lambda I - T) \leq 0$. The SVEP of T^* at λ entails that $q(\lambda I - T) < \infty$, see Remark 2.1, and by [1, Theorem 3.4] this implies that $\text{ind}(\lambda I - T) \geq 0$. Therefore, $\text{ind}(\lambda I - T) = 0$, i.e. $\alpha(\lambda I - T) = \beta(\lambda I - T) < \infty$ and consequently, again by [1, Theorem 3.4], $p(\lambda I - T) < \infty$, thus $\lambda \notin \sigma_b(T)$.

(ii) The inclusions $\sigma_{\text{wa}}(T^*) = \sigma_{\text{ws}}(T) \subseteq \sigma_w(T) \subseteq \sigma_b(T)$ hold for every $T \in L(X)$. Suppose that $\lambda \notin \sigma_{\text{wa}}(T^*)$. Then $\lambda I^* - T^* \in \Phi_+(X^*)$ with $\text{ind}(\lambda I^* - T^*) \leq 0$. By duality $\lambda I - T \in \Phi_-(X)$ and by Remark 2.1 the SVEP of T at λ entails that $p(\lambda I - T) < \infty$. By [1, Theorem 3.4] we have $\text{ind}(\lambda I - T) \leq 0$, thus $\text{ind}(\lambda I^* - T^*) = -\text{ind}(\lambda I - T) \geq 0$. Therefore, $\text{ind}(\lambda I^* - T^*) = \text{ind}(\lambda I - T) = 0$, and, again by [1, Theorem 3.4], we have $q(\lambda I - T) < \infty$, thus $\lambda \notin \sigma_b(T)$. \square

The following result has a crucial role in the sequel.

Theorem 2.7. Suppose that $T \in L(X)$. Then the following statements hold:

- (i) If T is polaroid and T^* has SVEP then property (w) holds for T .
- (ii) If T is polaroid and T has SVEP then property (w) holds for T^* .

Proof. (i) Note that by Corollary 2.45 of [1] we have $\sigma_a(T) = \sigma(T)$. Suppose first that $\text{iso } \sigma(T) = \emptyset$. Then $\pi_{00}(T) = \emptyset$. We show that also $\sigma_a(T) \setminus \sigma_{\text{wa}}(T)$ is empty. By Theorem 2.6 we have $\sigma_a(T) \setminus \sigma_{\text{wa}}(T) = \sigma(T) \setminus \sigma_b(T)$ and the last set is empty, since $\sigma(T)$ has no isolated points. Therefore, T satisfies property (w).

Consider the other case, $\text{iso } \sigma(T) \neq \emptyset$. Suppose that $\lambda \in \pi_{00}(T)$. Then λ is isolated in $\sigma(T)$ and hence, by the polaroid condition, λ is a pole of the resolvent of T , i.e. $p := p(\lambda I - T) = q(\lambda I - T) < \infty$. By assumption $\alpha(\lambda I - T) < \infty$, so by [1, Theorem 3.1] $\beta(\lambda I - T) < \infty$,

and hence $\lambda I - T \in \Phi(X)$. Therefore, by Theorem 2.6, $\lambda \in \sigma(T) \setminus \sigma_b(T) = \sigma(T) \setminus \sigma_{wa}(T)$. Conversely, if $\lambda \in \sigma_a(T) \setminus \sigma_{wa}(T) = \sigma(T) \setminus \sigma_b(T)$ then λ is an isolated point of $\sigma(T)$. Clearly, $0 < \alpha(\lambda I - T) < \infty$, so $\lambda \in \pi_{00}(T)$ and hence T satisfies property (w).

(ii) First note that since T has SVEP then $\sigma_a(T^*) = \sigma_s(T) = \sigma(T) = \sigma(T^*)$, see Corollary 2.45 of [1]. Suppose first that $\text{iso } \sigma(T) = \text{iso } \sigma(T^*) = \emptyset$. Then $\pi_{00}(T^*) = \emptyset$. By Theorem 2.6 we have $\sigma_a(T^*) \setminus \sigma_{wa}(T^*) = \sigma(T) \setminus \sigma_b(T) = \emptyset$, so T^* satisfies property (w).

Suppose that $\text{iso } \sigma(T) \neq \emptyset$ and let $\lambda \in \pi_{00}(T^*)$. Then λ is an isolated point of $\sigma(T^*) = \sigma(T)$, hence a pole of the resolvent of T^* , since T^* is polaroid by Theorem 2.5. By assumption $\alpha(\lambda I^* - T^*)^p < \infty$ and since the ascent and the descent of $\lambda I^* - T^*$ are both finite it then follows by [1, Theorem 3.1] that $\beta(\lambda I - T) = \alpha(\lambda I - T) < \infty$, so $\lambda I^* - T^*$ is Browder and hence also $\lambda I - T$ is Browder. Therefore, $\lambda \in \sigma(T) \setminus \sigma_b(T)$ and by Theorem 2.6 it then follows that $\lambda \in \sigma_a(T^*) \setminus \sigma_{wa}(T^*)$.

Conversely, if $\lambda \in \sigma_a(T^*) \setminus \sigma_{wa}(T^*) = \sigma(T) \setminus \sigma_b(T)$, then λ is an isolated point of the spectrum of $\sigma(T) = \sigma(T^*)$, $\lambda I - T \in B(X)$, or equivalently $\lambda I^* - T^* \in B(X^*)$. Since $\alpha(\lambda I^* - T^*) = \beta(\lambda I^* - T^*)$ we then have $\alpha(\lambda I^* - T^*) > 0$ (otherwise $\lambda \notin \sigma(T^*)$). Clearly, $\alpha(\lambda I^* - T^*) < \infty$, since by assumption $\lambda I^* - T^* \in W_+(X^*)$, so that $\lambda \in \pi_{00}(T^*)$. Thus T^* satisfies property (w). \square

The following example shows that in the statements (i) of Theorem 2.7 the assumption that T^* has SVEP cannot be replaced by the assumption that T has SVEP.

Example 2.8. Denote by R the unilateral right shift on $\ell^2(\mathbb{N})$ and define

$$U(x_1, x_2, \dots) := (0, x_2, x_3, \dots) \quad \text{for all } (x_n) \in \ell^2(\mathbb{N}).$$

Clearly, U is a quasi-nilpotent operator. Let $T := R \oplus U$. We have $\sigma(T) = \mathbf{D}$, \mathbf{D} the closed unit disc of \mathbb{C} , so $\text{iso } \sigma(T) = \pi_{00}(T) = \emptyset$ and hence T is polaroid. Moreover, $\sigma_a(T) = \partial\sigma(T) \cup \{0\}$. By (4) T has SVEP at 0, as well as at the points $\lambda \notin \sigma_a(T)$. Since T has SVEP at all points $\lambda \in \partial\sigma(T)$ it then follows that T has SVEP. Finally, $\sigma_{wa}(T) = \partial\sigma(T)$ so $\sigma(T) \setminus \sigma_{wa}(T) = \{0\} \neq \pi_{00}(T) = \emptyset$, thus T does not satisfy property (w).

Analogously, in the statements (ii) of Theorem 2.7 the assumption that T has SVEP cannot be replaced by the assumption that T^* has SVEP.

Example 2.9. Let us consider the left shift $L \in L(\ell^2(\mathbb{N}))$, and let U' be the adjoint of the quasi-nilpotent operator U defined in Example 2.8. We have $L' = R$, R the unilateral right shift. If we define $S := L \oplus U'$ then, as observed in Example 2.8 $S' = R \oplus U$ has SVEP (and hence the dual S^* has SVEP). From Example 2.8 we also know that $\sigma(S) = \overline{\sigma(S')} = \mathbf{D}$, so S is polaroid and S' has not property (w), or equivalently by Theorem 2.4, S^* has not property (w).

In the sequel by $H_0(T)$ we shall denote the *quasi-nilpotent part* of T , defined as

$$H_0(T) := \{x \in X : \lim_{n \rightarrow \infty} \|T^n x\|^{1/n} = 0\}.$$

It is easily seen that $\ker(T^m) \subseteq H_0(T)$ for every $m \in \mathbb{N}$. Another important subspace in local spectral theory is given by *analytic core* of T , defined as the set $K(T)$ of all $x \in X$ such that there exists a sequence $(u_n) \subset X$ and $\delta > 0$ for which $x = u_0$, and $Tu_{n+1} = u_n$ and $\|u_n\| \leq \delta$.

$\delta^n \|x\|$ for every $n \in \mathbb{N}$. It easily follows, from the definition, that $K(T)$ is a linear subspace of X and that $T(K(T)) = K(T)$. Note that, (see [4]),

$$H_0(\lambda I - T) \text{ closed} \Rightarrow T \text{ has SVEP at } \lambda, \quad (7)$$

and this implication is an equivalence whenever $\lambda I - T \in \Phi_{\pm}(X)$, see [4].

Theorem 2.10. *If $T \in L(X)$ is polaroid and $N \in L(X)$ is nilpotent such that $TN = NT$. Then $T + N$ is polaroid.*

Proof. Note first that the quasi-nilpotent part of every operator $S \in L(X)$ is invariant under nilpotent perturbations. In fact, let $x \in H_0(S)$ and suppose that $N^{\nu} = 0$ for some $\nu \in \mathbb{N}$. If $U := \sum_{k=0}^{\nu-1} c_{\nu,k} S^{\nu-1-k} N^k$, with suitable binomial coefficients $c_{\nu,k}$, then it is easily seen that $(S + N)^{\nu} = SU$, from which the following estimate follows

$$\|(S + N)^{\nu} x\|^{\frac{1}{\nu}} \leq \|S^{\nu} x\|^{\frac{1}{\nu}} \|U^{\nu} x\|^{\frac{1}{\nu}}.$$

From this estimate we deduce that $\lim_{n \rightarrow +\infty} \|(S + N)^{\nu n} x\|^{\frac{1}{n}} = 0$, i.e. $x \in H_0(S + N)$, and hence $H_0(S) \subseteq H_0(S + N)$. The reverse inclusion follows by symmetry:

$$H_0(S + N) \subseteq H_0((S + N) - N) = H_0(S).$$

To show that $T + N$ is polaroid, let $\lambda \in \text{iso } \sigma(T + N)$. It is well-known that the spectrum of an operator is invariant under nilpotent perturbations, so λ is an isolated point in $\sigma(T)$, hence a pole of the resolvent of T . Let $p := p(\lambda I - T) = q(\lambda I - T)$. By the first part of the proof and by Theorem 3.74 of [1] we have

$$H_0(\lambda I - T + N) = H_0(\lambda I - T) = \ker(\lambda I - T)^p.$$

Set $m := p\nu$. We show that

$$H_0(\lambda I - T + N) = \ker(\lambda I - T + N)^m. \quad (8)$$

The inclusion $\ker(\lambda I - T + N)^n \subseteq H_0(\lambda I - T + N)$ is true for all operators and for all $n \in \mathbb{N}$. Suppose that $x \in H_0(\lambda I - T + N) = \ker(\lambda I - T)^p$. Then we can write

$$(\lambda I - T + N)^m x = \sum_{i=m-p+1}^m [\mu_{i,m}(\lambda I - T)^{m-i} N^{i-\nu}] N^{\nu} x = 0,$$

with suitable binomial coefficients $\mu_{i,m}$, thus

$$H_0(\lambda I - T + N) = \ker(\lambda I - T)^p \subseteq \ker(\lambda I - T + N)^m$$

and hence (8) is proved.

We show now that $T + N$ is polaroid. Let $\lambda \in \text{iso } \sigma(T + N)$. Then, by Theorem 3.74 of [1], we have

$$X = H_0(\lambda I - T + N) \oplus K(\lambda I - T + N) = \ker(\lambda I - T + N)^m \oplus K(\lambda I - T + N),$$

so that

$$(\lambda I - T + N)^m(X) = (\lambda I - T + N)^m(K(\lambda I - T + N)) = K(\lambda I - T + N).$$

Therefore, $X = \ker(\lambda I - T + N)^m \oplus (\lambda I - T + N)^m(X)$ and this decomposition implies, by Theorem 3.6 of [1], that $\lambda I - T + N$ has both finite ascent and descent, or equivalently that λ is a pole of the resolvent of $T + N$. \square

Theorem 2.11. Suppose that $T \in L(X)$ is polaroid, $N \in L(X)$ a nilpotent operator commuting with T .

- (i) If T has SVEP then $T^* + N^*$ satisfies property (w), or equivalently a -Weyl's theorem holds for $T^* + N^*$.
- (ii) If T^* has SVEP then $T + N$ satisfies property (w), or equivalently a -Weyl's theorem holds for $T + N$.

Proof. (i) If T has SVEP then $T + N$ has SVEP, see Corollary 2.12 of [1]. Moreover, by Lemma 2.10 $T + N$ is polaroid. By Theorem 2.7 it then follows that property (w) holds for $T^* + N^*$, or equivalently, since $T + N$ has SVEP, a -Weyl's theorem holds for $T^* + N^*$.

(ii) If T is polaroid then by Theorem 2.5 T^* is polaroid. Clearly, N^* is nilpotent, since $(N^*)^n = (N^n)^* = 0$ for some $n \in \mathbb{N}$. Therefore $T^* + N^*$ is polaroid, by Theorem 2.10. Since $T^* + N^*$ has SVEP, by Corollary 2.12 of [1], it then follows, by Theorem 2.7, that $T + N$ satisfies property (w), or equivalently a -Weyl's theorem. \square

We can say much more than Theorem 2.11. Denote by $\mathcal{H}(\sigma(T))$ the set of all analytic functions defined on a neighborhood \mathcal{U} of $\sigma(T)$, let $f(T)$ be defined by means of the classical functional calculus:

$$f(T) := \frac{1}{2\pi i} \int_{\Gamma} f(\lambda)(\lambda I - T)^{-1} d\lambda,$$

where Γ is a contour in \mathcal{U} that surrounds $\sigma(T)$.

Theorem 2.12. Suppose that T is polaroid and $N \in L(X)$ a nilpotent operator commuting with T . If T^* has SVEP and $f \in H(\sigma(T))$ then property (w) holds for $f(T) + N$, or equivalently a -Weyl's theorem holds for $f(T) + N$.

Proof. By Theorem 2.7, T satisfies property (w), or equivalently, by Theorem 2.2, T satisfies Weyl's theorem. The SVEP for T^* implies that $\sigma(T) = \sigma_a(T)$, see [1, Corollary 2.45], so every isolated point of $\sigma_a(T)$ is a pole of the resolvent of T . By [6, Theorem 2.22] then $f(T)$ satisfies property (w). Finally, by [3, Theorem 2.8] $f(T) + N$ satisfies property (w). Since $f(T^*) = f(T)^*$ has SVEP, see Theorem 2.40 of [1], by Theorem 2.2 it then follows that property (w) and a -Weyl's theorem for $f(T) + N$ are equivalent. \square

Remark 2.13. It is somewhat meaningful to ask what we can say about the operators $f(T + N)$, always under the assumptions of Theorem 2.12. Now, if T is polaroid then $T + N$ is polaroid, by Theorem 2.10. Moreover, by $T^* + N^* = (T + N)^*$ has SVEP by Corollary 2.12 of [1]. Now, as it has been observed in the proof of Theorem 2.12, if $S := T + N$ is polaroid and $S^* = (T + N)^*$ has SVEP then S is a -polaroid. Hence by [6, Theorem 2.22] $f(T + N)$ satisfies property (w) for every $f \in \mathcal{H}(\sigma(T))$.

A bounded operator $T \in L(X)$ is said to have property $H(p)$ if for all $\lambda \in \mathbb{C}$ there exists a $p := p(\lambda) \in \mathbb{N}$ such that:

$$H_0(\lambda I - T) = \ker(\lambda I - T)^p.$$

This class of operators has been introduced in [18] and for the constant function $p(\lambda) := 1$ has been also studied in [7]. Clearly, from the implication (7), if T has property $H(p)$ then T has SVEP. Moreover T is polaroid, see [2, Lemma 3.3].

It should be noted that property $H(p)$ holds for a relevant number of Hilbert space operators. In [18, Example 3] Oudghiri observed that every *generalized scalar operator* and every *subscalar operator* T (i.e. T is similar to a restriction of a generalized scalar operator to one of its closed invariant subspaces, we refer to [15] for definitions and results) has property $H(p)$, see [15] for definitions and properties. Consequently, property $H(p)$ is satisfied by p -hyponormal operators and *log*-hyponormal operators [16, Corollary 2], w -hyponormal operators [17], M -hyponormal operators [15, Proposition 2.4.9], and totally paranormal operators [7]. Also totally $*$ -paranormal operators have property $H(1)$ [13]. Property $H(p)$ for $p := p(\lambda) = 1$ also holds for every multiplier $T \in M(A)$ of a commutative semi-simple Banach algebra A , and in particular for every convolution operator on $L^1(G)$, where G is a compact Abelian group [7].

A bounded operator $T \in L(X)$ on a Banach space X is said to be *paranormal* if

$$\|Tx\|^2 \leq \|T^2x\|\|x\| \quad \text{holds for all } x \in X.$$

The class of paranormal operators properly contains a relevant number of Hilbert space operators, among them p -hyponormal operators, *log*-hyponormal operators, and the class of operators satisfying the condition $|T|^2 \leq |T^2|$, see [12] and [10]. Note that, in general, paranormal operators do not satisfy property $H(p)$, see [5] for a counter-example. A bounded operator $T \in L(X)$ is said to be *algebraically paranormal* if there exists a non-trivial polynomial h such that $h(T)$ is paranormal.

Every paranormal operator on a Hilbert space has SVEP. This may be easily seen as follows: if $\lambda \neq 0$ and $\lambda \neq \mu$ then, by Theorem 2.6 of [8], we have

$$\|x + y\| \geq \|y\| \quad \text{for all } x \in \ker(\mu I - T) \text{ and } y \in \ker(\lambda I - T). \quad (9)$$

It then follows that if U is an open disc and $f : U \rightarrow X$ is an analytic function such that $0 \neq f(z) \in \ker(zI - T)$ for all $z \in U$, then f fails to be continuous at every $0 \neq \lambda \in U$. By Theorem 2.40 of [1] it then follows that every algebraically paranormal satisfies SVEP.

Recall that a bounded operator T is said to be *algebraic* if there exists a non-trivial polynomial h such that $h(T) = 0$. From the spectral mapping theorem it easily follows that the spectrum of an algebraic operator is a finite set. A nilpotent operator is a trivial example of an algebraic operator. Also finite rank operators K are algebraic; more generally, if K^n is a finite rank operator for some $n \in \mathbb{N}$ then K is algebraic. Clearly, if T is algebraic then its dual T^* is algebraic, as well as T' in the case of Hilbert space operators.

Theorem 2.14. *Suppose that $T \in L(X)$ and $K \in L(X)$ is an algebraic operator commuting with T .*

- (i) *If T has SVEP then $T + K$ has SVEP.*
- (ii) *If T is algebraically paranormal, or $T \in H(p)$, then $T + K$ is polaroid.*

Proof. (i) Let $\sigma(K) = \{\mu_1, \dots, \mu_n\}$. Denote by P_i the spectral projection associated with K and with the spectral set $\{\mu_i\}$. If $Y_i := P_i(H)$ and $Z_i := \ker P_i$, then $H = Y_i \oplus Z_i$, the closed subspaces Y_i and Z_i are invariant under T and K , and $\sigma(K|Y_i) = \{\mu_i\}$. Define $K_i := K|Y_i$ and $T_i := T|Y_i$. Clearly, the restrictions T_i and K_i commute for every $i = 1, 2, \dots, n$. Let h be a polynomial such that $h(K) = 0$. Then $h(K_i) = h(K)|Y_i = 0$, and the equalities

$$\{0\} = \sigma(h(K_i)) = h(\sigma(K_i)) = h(\{\mu_i\})$$

entail that $h(\mu_i) = 0$. Write

$$h(\mu) = (\mu_i - \mu)^v q(\mu) \quad \text{with } q(\mu_i) \neq 0.$$

Then $0 = h(K_i) = (\mu_i I - K_i)^v q(K_i)$ where the operators $q(K_i)$ is invertible. Therefore $(\mu_i I - K_i)^v = 0$, hence the operators $N_i := \mu_i I - K_i$ are nilpotent for all $i = 1, 2, \dots, n$. Note that

$$T_i + K_i = (\mu_i I + T_i) + (K_i - \mu_i I) = \mu_i I + T_i - N_i. \quad (10)$$

Since SVEP is inherited by restrictions to closed invariant subspaces, T_i has SVEP and hence, by [1, Corollary 2.12] also $T_i + K_i = \mu_i I + T_i - N_i$ has SVEP for all $i = 1, 2, \dots, n$. By Theorem 2.9 of [1] it then follows that

$$T + K = \bigoplus_{i=1}^n T_i + K_i$$

has SVEP.

(ii) Suppose that T is algebraically paranormal. Let $\lambda \in \text{iso } \sigma(T + K)$. Since $\sigma(T + K) = \bigcup_{i=1}^n \sigma(T_i + K_i)$ then $\lambda \in \text{iso } \sigma(T_j + K_j)$ for some positive integer $1 \leq j \leq n$ and hence $\lambda - \mu_j \in \text{iso } \sigma(T_j + K_j - \mu_j I)$. The restriction to a closed invariant subspace of an algebraically paranormal is algebraically paranormal, so T_j is polaroid. Since, as observed before, $\mu_j I - K_j$ is nilpotent then, by Theorem 2.10, also $T_j + K_j - \mu_j I$ is polaroid. Therefore $\lambda - \mu_j$ is a pole of the resolvent of $T_j + K_j - \mu_j I$, so there exists by Theorem 3.74 of [1] a positive integer p_j such that

$$H_0[(\lambda - \mu_j)I - (T_j + K_j - \mu_j I)] = H_0(\lambda I - (T_j + K_j)) = \ker(\lambda I - (T_j + K_j))^{p_j}.$$

Therefore, taking in to account that $H_0(\lambda I - (T_i + K_i)) = \{0\}$ if $\lambda \notin \sigma(T_i + K_i)$, we have

$$\begin{aligned} H_0(\lambda I - (T + K)) &= \bigoplus_{i=1}^n H_0(\lambda I - (T_i + K_i)) = \bigoplus_{i=1}^n \ker(\lambda I - (T_i + K_i))^{p_i} \\ &= \ker(\lambda I - (T + K))^p, \end{aligned}$$

where $p := \max\{p_1, p_2, \dots, p_n\}$. Arguing as in the proof of Theorem 2.10 it then follows that λ is a pole of the resolvent of $T + K$. The assertion concerning $H(p)$ -operators may be proved in a similar way, just observe that if $T \in H(p)$ then every restriction of T to a closed invariant subspaces has property $H(p)$ and hence is polaroid, see [18]. \square

In [5] it has been proved that if $T \in L(H)$ is paranormal, K is an algebraic operator commuting with T , then Weyl's theorem holds for $T + K$. Something more can be said.

Theorem 2.15. *Suppose that $T \in L(H)$, H a Hilbert space and $K \in L(H)$ is an algebraic operator commuting with T .*

- (i) *If T is algebraically paranormal then property (w) holds for $T' + K'$.*
- (ii) *If T' is algebraically paranormal then property (w) holds for $T + K$.*

Proof. (i) If T is algebraically paranormal then T has SVEP and hence $T + K$ has SVEP. Moreover, T is polaroid so also $T + K$ is polaroid. By Theorem 2.7, part (i), then property (w) holds for $T^* + K^*$ and this is equivalent, by Lemma 2.4, to saying that property (w) holds for $T' + K'$.

(ii) If T' is algebraically paranormal then T' has SVEP and hence $T' + K'$ has SVEP, equivalently $T^* + K^*$ has SVEP. Moreover, $T' + K'$ is polaroid, so, by Theorem 2.5, $T + K$ is polaroid. By Theorem 2.7 it then follows that property (w) holds for $T + K$. \square

In [19] it has been proved that if $T \in H(p)$, K is an algebraic operator commuting with T , then Weyl's theorem holds for $T + K$. A similar result of that of Theorem 2.15 holds for operators $T \in H(p)$.

Theorem 2.16. *Suppose that $T \in L(X)$ and $K \in L(X)$ is an algebraic operator commuting with T .*

- (i) *If $T \in H(p)$ then property (w) holds for $T^* + K^*$.*
- (ii) *If $T^* \in H(p)$ then property (w) holds for $T + K$.*

Proof. Proceed as in the proof of Theorem 2.15. \square

Theorem 2.15 and Theorem 2.16 may be somewhat extended as follows: Recall that a subspace M of X is said to be *orthogonal* (in the sense of Birkhoff) to an other subspace N of X , $M \perp N$ if $\|x\| \leq \|x + y\|$ for all $x \in M$, $y \in N$. This asymmetric definition coincides with the usual concept of orthogonality in the case that X is a Hilbert space. M and N are said to be *mutually orthogonal*, $M \perp_m N$ if $M \perp N$ and $N \perp M$. From (9) it then follows that paranormal operators satisfy the *orthogonality condition*

$$\ker(\mu I - T) \perp_m \ker(\lambda I - T) \quad \text{for all } \lambda \neq \mu, \quad \lambda \neq 0, \quad (11)$$

and the same argument used for paranormal operators shows that this condition entails SVEP for T . Recall that $T \in L(X)$ is said to be *normaloid* if $\|T\| = r(T)$, $r(T)$ the spectral radius of T . A bounded operator T is said *completely hereditarily normaloid*, $T \in \mathcal{CHN}$, if either every part, and (also) every invertible part, of T is normaloid, or for all $\lambda \in \mathbb{C}$ every part of $\lambda I - T$ is normaloid.

Theorem 2.17. *Suppose that $T \in \mathcal{CHN}$ satisfies the orthogonality condition (11). If $K \in L(X)$ is algebraic and commutes with T then $T^* + K^*$ satisfies property (w).*

Proof. Every $T \in \mathcal{CHN}$ is polaroid, so $T + K$ is polaroid. The orthogonal condition (11) entails that T has SVEP, hence also $T + K$ has SVEP, by Lemma 2.14. The result then follows from Theorem 2.7, part (ii). \square

Acknowledgments

The authors thank the referee for his several suggestions which have greatly contributed to improve the final form of this article.

References

- [1] P. Aiena, *Fredholm and Local Spectral Theory, with Application to Multipliers*, Kluwer Academic Publishers, 2004.
- [2] P. Aiena, Classes of operators satisfying a -Weyl's theorem, *Studia Math.* 169 (2005) 105–122.
- [3] P. Aiena, M.T. Biondi, Property (w) and perturbations, *J. Math. Anal. Appl.* 336 (2007) 683–692.
- [4] P. Aiena, M.L. Colasante, M. Gonzalez, Operators which have a closed quasi-nilpotent part, *Proc. Amer. Math. Soc.* 130 (9) (2002) 2701–2710.
- [5] P. Aiena, J.R. Guillen, Weyl's theorem for perturbations of paranormal operators, *Proc. Amer. Math. Anal. Soc.* 35 (2007) 2433–2442.
- [6] P. Aiena, P. Peña, A variation on Weyl's theorem, *J. Math. Anal. Appl.* 324 (2006) 566–579.
- [7] P. Aiena, F. Villafañe, Weyl's theorem for some classes of operators, *Integral Equations Operator Theory* 53 (2005) 453–466.
- [8] N.N. Chourasia, P.B. Ramanujan, Paranormal operators on Banach spaces, *Bull. Austral. Math. Soc.* 21 (1980) 161–168.
- [9] L.A. Coburn, Weyl's theorem for nonnormal operators, *Michigan Math. J.* 20 (1970) 529–544.
- [10] B.P. Duggal, Roots of contraction with Hilbert–Schmidt defect operators and C_0 completely non unitary part, *Comm. Math. Ann. Soc. Polonae* 36 (1996) 85–106.
- [11] B.P. Duggal, Polaroid operators satisfying Weyl's theorem, *Linear Algebra Appl.* 414 (2006) 271–277.
- [12] T. Furuta, M. Ito, T. Yamazaki, A subclass of paranormal operators including class of log-hyponormal and several related classes, *Sci. Math.* 1 (1998) 389–403.
- [13] Y.M. Han, An-Hyun Kim, A note on $*$ -paranormal operators, *Integral Equations Operator Theory* 49 (2004) 435–444.
- [14] H. Heuser, *Functional Analysis*, Marcel Dekker, New York, 1982.
- [15] K.B. Laursen, M.M. Neumann, *Introduction to Local Spectral Theory*, Clarendon Press, Oxford, 2000.
- [16] C. Lin, Y. Ruan, Z. Yan, p -hyponormal operators are subscalar, *Proc. Amer. Math. Soc.* 131 (9) (2003) 2753–2759.
- [17] C. Lin, Y. Ruan, Z. Yan, w -hyponormal operators are subscalar, *Integral Equations Operator Theory* 50 (2004) 165–168.
- [18] M. Oudghiri, Weyl's and Browder's theorem for operators satisfying the SVEP, *Studia Math.* 163 (1) (2004) 85–101.
- [19] M. Oudghiri, Weyl's theorem and perturbations, *Integral Equations Operator Theory* 53 (2005) 535–545.
- [20] V. Rakočević, On a class of operators, *Mat. Vesnik* 37 (1985) 423–426.
- [21] V. Rakočević, Operators obeying a -Weyl's theorem, *Rev. Roumaine Math. Pures Appl.* 34 (10) (1989) 915–919.